

# A Determination of the Weak Phase $\gamma$ from Color-allowed $B_u^\pm \rightarrow DK^\pm$ Decays

Zhi-zhong Xing<sup>1</sup>

*Sektion Physik, Universität München, Theresienstrasse 37A, 80333 München, Germany*

## Abstract

We show that it is possible to determine the weak phase  $\gamma \equiv \arg(-V_{ub}^* V_{ud}/V_{cb}^* V_{cd})$  of the Cabibbo-Kobayashi-Maskawa flavor mixing matrix only from the measurement of the color-allowed  $B_u^\pm \rightarrow DK^\pm$  decay rates. The uncertainty of this method, arising mainly from the factorization approximation for two tree-level spectator quark transitions, may be well controlled.

PACS number(s): 12.15.Hh, 12.15.Ji, 13.25.Hw, 14.40.Nd

---

<sup>1</sup>Electronic address: xing@hep.physik.uni-muenchen.de

The first observation of the Cabibbo-suppressed but color-allowed transitions  $B_u^+ \rightarrow \bar{D}^0 K^+$  and  $B_u^- \rightarrow D^0 K^-$ , which involve the quark subprocesses  $b \rightarrow c\bar{u}s$  and  $\bar{b} \rightarrow \bar{c}u\bar{s}$  respectively, has been reported by the CLEO Collaboration [1]. These two decay modes, together with  $B_u^+ \rightarrow D^0 K^+$ ,  $B_u^+ \rightarrow D_{1,2} K^+$  and their charge-conjugate counterparts <sup>2</sup>, can in principle be used to determine the weak phase  $\gamma \equiv \arg(-V_{ub}^* V_{ud}/V_{cb}^* V_{cd})$  of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [2, 3] <sup>3</sup>. In practice, however, such a method may have an essential problem, arising from the difficulty in measuring the color-suppressed transitions  $B_u^+ \rightarrow D^0 K^+$  and  $B_u^- \rightarrow \bar{D}^0 K^-$ . To get around this problem, Atwood, Dunietz and Soni have proposed the measurement of  $B_u^+ \rightarrow D^0 K^+ \rightarrow (K^-\pi^+)_{D^0} K^+$ ,  $B_u^+ \rightarrow \bar{D}^0 K^+ \rightarrow (K^-\pi^+)_{\bar{D}^0} K^+$  and their charge-conjugate channels for the extraction of  $\gamma$  [5]. Furthermore Soffer has shown how to remove some uncertainties from the methods mentioned above and increase their sensitivity by measuring  $CP$ -conserving phases at a  $\tau$ -charm factory [6].

The interesting question remains if it is possible to constrain  $\gamma$  only from observation of the color-allowed  $B_u^\pm \rightarrow DK^\pm$  decay rates. Gronau has recently discussed this possibility by proposing a few promising measurables, such as the charge-averaged ratios for  $B_u^\pm$  decays into  $D$ -meson  $CP$  and flavor states [7]:

$$R_i \equiv 2 \frac{\Gamma(B_u^+ \rightarrow D_i K^+) + \Gamma(B_u^- \rightarrow D_i K^-)}{\Gamma(B_u^+ \rightarrow \bar{D}^0 K^+) + \Gamma(B_u^- \rightarrow D^0 K^-)} \quad (1)$$

with  $i = 1$  or  $2$  (the factor 2 on the right-hand side of Eq. (1) is just taken to normalize  $R_i$  to a value close to one). It is easy to show that there exist two inequalities,  $\sin^2 \gamma \leq R_{1,2}$ ; and one of the two ratios  $R_1$  and  $R_2$  must be smaller than one except that the value of  $\gamma$  happens to lie in a narrow band around  $\pi/2$ . Thus the measurement of  $R_{1,2}$  may in most cases provide a useful constraint on  $\gamma$ .

The purpose of this note is to show that  $\gamma$  can indeed be *determined* from  $R_1$  and  $R_2$  through the formula

$$\cos \gamma = \frac{\kappa}{2} \frac{R_1 - R_2}{R_1 + R_2 - 2} \quad , \quad (2)$$

where  $\kappa \approx 0.077$  is a coefficient related to the ratio of the color-suppressed and color-allowed decay amplitudes:

$$r \equiv \left| \frac{A(B_u^+ \rightarrow D^0 K^+)}{A(B_u^+ \rightarrow \bar{D}^0 K^+)} \right|$$

---

<sup>2</sup>where  $D_1$  and  $D_2$  denote the  $CP$ -even and  $CP$ -odd states of  $D^0$  and  $\bar{D}^0$  mesons, respectively.

<sup>3</sup>Possible new physics in  $D^0$ - $\bar{D}^0$  mixing could affect those methods proposed in Refs. [2, 3]. See Ref. [4] for some detailed discussions.

$$= \left| \frac{A(B_u^- \rightarrow \bar{D}^0 K^-)}{A(B_u^- \rightarrow D^0 K^-)} \right|. \quad (3)$$

The value of  $\kappa$  is obtained by means of the isospin symmetry and the factorization approximation. Since the factorization approximation made here involves only tree-level spectator quark diagrams, the uncertainty associated with  $\kappa$  may be well controlled. In fact, the reported branching ratio  $\mathcal{B}(B_u^+ \rightarrow \bar{D}^0 K^+) = (0.257 \pm 0.065 \pm 0.032) \times 10^{-3}$  [1] is in good agreement with the prediction from the factorization scheme [8]. Therefore the measurement of the color-allowed  $B_u^\pm \rightarrow DK^\pm$  decay rates should allow a determination of the weak phase  $\gamma$  from Eq. (2) to an acceptable degree of accuracy in the near future.

To derive the formula in Eq. (2) and calculate the parameter  $\kappa$ , we begin with the effective weak Hamiltonians responsible for  $b \rightarrow c\bar{u}s$ ,  $b \rightarrow u\bar{c}s$  and their charge-conjugate transitions:

$$\begin{aligned} \mathcal{H}_{\text{eff}}^{(1)} &= \frac{G_F}{\sqrt{2}} V_{cb} V_{us}^* \left[ c_1 (\bar{s}u)_{V-A} (\bar{c}b)_{V-A} + c_2 (\bar{c}u)_{V-A} (\bar{s}b)_{V-A} \right] + \text{h.c.}, \\ \mathcal{H}_{\text{eff}}^{(2)} &= \frac{G_F}{\sqrt{2}} V_{ub} V_{cs}^* \left[ c_1 (\bar{s}c)_{V-A} (\bar{u}b)_{V-A} + c_2 (\bar{u}c)_{V-A} (\bar{s}b)_{V-A} \right] + \text{h.c.}, \end{aligned} \quad (4)$$

where  $c_1$  and  $c_2$  are QCD correction coefficients. To calculate the hadronic matrix elements  $\langle DK | \mathcal{H}_{\text{eff}} | B \rangle$ , one has to make some approximations. Here we use the factorization approximation, which factorizes each four-quark operator matrix element into a product of two current matrix elements. Wherever there is a color mismatch between the quark operator and the external state in this approach, a phenomenological parameter  $\xi$  is introduced and the factorized matrix element becomes related to one of the following two coefficients [9]:

$$a_1 \equiv c_1 + \xi c_2, \quad a_2 \equiv c_2 + \xi c_1. \quad (5)$$

The color-suppression factor  $\xi$  is naively expected to be  $1/3$ , corresponding to exact vacuum saturation. Both  $a_1$  and  $a_2$  have been determined from experimental data [10]. The explicit knowledge of  $\xi$ ,  $c_1$  and  $c_2$  will be irrelevant in our subsequent analysis.

When applying the effective Hamiltonians and the factorization approximation to  $B_u^\pm \rightarrow DK^\pm$  decays, one has to take possible final-state interactions into account. Since both  $D$  and  $K$  are isospin  $1/2$  particles, the state  $DK$  can be either  $I = 0$  or  $I = 1$ . An isospin analysis made by Deshpande and Dib [11] gives <sup>4</sup>

$$A(B_u^- \rightarrow \bar{D}^0 K^-) = \frac{G_F}{\sqrt{2}} (V_{ub} V_{cs}^*) \left( \frac{X}{2} e^{i\phi_0} + \frac{X}{2} e^{i\phi_1} \right),$$

---

<sup>4</sup>The factorized matrix element  $\langle \bar{D}^0 K^- | (\bar{s}c)_{V-A} | 0 \rangle \langle 0 | (\bar{u}b)_{V-A} | B_u^- \rangle$ , corresponding to an annihilation process, is expected to be formfactor-suppressed [12, 13] and has been neglected here.

$$A(B_u^- \rightarrow D^0 K^-) = \frac{G_F}{\sqrt{2}} (V_{cb} V_{us}^*) (X + Y) e^{i\phi_1}, \quad (6)$$

where  $\phi_0$  and  $\phi_1$  are the strong phase parameters,  $X$  and  $Y$  are the factorized hadronic matrix elements:

$$\begin{aligned} X &= a_2 \langle \bar{D}^0 | (\bar{u}c)_{V-A} | 0 \rangle \langle K^- | (\bar{s}b)_{V-A} | B_u^- \rangle \\ Y &= a_1 \langle K^- | (\bar{s}u)_{V-A} | 0 \rangle \langle D^0 | (\bar{c}b)_{V-A} | B_u^- \rangle. \end{aligned} \quad (7)$$

Clearly the weak phase difference between  $A(B_u^- \rightarrow \bar{D}^0 K^-)$  and  $A(B_u^- \rightarrow D^0 K^-)$  amounts to  $-\gamma$  to an excellent degree of accuracy in the standard model [4]. The strong phase difference between these two amplitudes, denoted by  $\delta$  as the notation in Ref. [7], is equal to  $(\phi_0 - \phi_1)/2$ . By convention, we take  $(\phi_1 - \phi_0) \in [-\pi, +\pi]$ . Then  $\cos \delta \geq 0$  holds.

In Ref. [7] the ratio  $r$  and the strong phase difference  $\delta$  are formally taken as two independent parameters. In our factorization approach, however,  $r$  depends on  $\delta$  through the relationship  $r = \kappa \cos \delta$  with

$$\kappa = \left| \frac{V_{ub} V_{cs}^*}{V_{cb} V_{us}^*} \right| \left| \frac{X}{X + Y} \right|, \quad (8)$$

derived from Eqs. (3) and (6). The point is simply that the amplitude of  $B_u^- \rightarrow \bar{D}^0 K^-$ , after its CKM coefficient is factored out, contains two isospin components with equal magnitude but different phases. The result of  $R_{1,2}$  obtained by Gronau [7] can be reproduced as follows:

$$R_{1,2} = 1 + r^2 \pm 2r \cos \delta \cos \gamma. \quad (9)$$

By use of  $r = \kappa \cos \delta$ , we arrive at

$$R_{1,2} = 1 + \kappa \cos^2 \delta (\kappa \pm 2 \cos \gamma). \quad (10)$$

It is easy to obtain two inequalities,  $\sin^2 \gamma \leq R_{1,2}$ , from either Eq. (9) or Eq. (10). Note that there exists a narrow band around  $\gamma = \pi/2$ , which makes both  $R_1$  and  $R_2$  equal to or larger than one. With the help of Eq. (10), we find that the necessary condition for  $R_{1,2} \geq 1$  is

$$-\frac{\kappa}{2} \leq \cos \gamma \leq +\frac{\kappa}{2}. \quad (11)$$

Beyond this band of  $\gamma$ , one of the two ratios  $R_1$  and  $R_2$  must be smaller than one, thus one may get the constraint  $\sin^2 \gamma \leq R_1 < 1$  or  $\sin^2 \gamma \leq R_2 < 1$ .

Note that the weak phase  $\gamma$  can indeed be determined from  $R_1$  and  $R_2$ , provided the value of  $\kappa$  is known. From Eq. (10), we straightforwardly obtain

$$\cos \gamma = \frac{\kappa}{2} \frac{R_1 - R_2}{R_1 + R_2 - 2}. \quad (12)$$

This instructive result has been listed in Eq. (2). To evaluate  $\kappa$ , we express the hadronic matrix elements  $X$  and  $Y$  in terms of relevant decay constants and formfactors. The ratio  $X/Y$  turns out to be

$$\frac{X}{Y} = \frac{a_2}{a_1} \frac{m_B^2 - m_K^2}{m_B^2 - m_D^2} \frac{f_D}{f_K} \frac{F_0^{BK}(m_D^2)}{F_0^{BD}(m_K^2)}. \quad (13)$$

The main error bar of  $X/Y$  comes from the unknown decay constant  $f_D$ . Here we typically take  $f_D = 220$  MeV. We also input  $f_K = 160$  MeV,  $a_2/a_1 = 0.25$  [10],  $F_0^{BK}(0) = 0.38$ , and  $F_0^{BD}(0) = 0.69$  [9]. The uncertainties of the chosen values for  $F_0^{BK}(0)$  and  $F_0^{BD}(0)$  are expected to be within 15% [14]. By use of a simple monopole model for formfactors [9], we obtain  $F_0^{BK}(m_D^2) \approx 0.43$  and  $F_0^{BD}(m_K^2) \approx 0.70$ . Then we arrive at  $X/Y \approx 0.24$ . With the input  $|(V_{ub}V_{cs}^*)/(V_{cb}V_{us}^*)| = 0.4$  [15], we finally get  $\kappa \approx 0.077$ . This implies that the value of  $r = \kappa \cos \delta$  is smaller than the naive expectation  $r \approx 0.1$ , obtained in Ref. [7] with the assumption  $\delta = 0$  and  $X/(X + Y) \approx a_2/a_1$ . Since the error associated with the CKM factor is only 25% or so [15] and those associated with the formfactors can be partly cancelled in the ratio  $X/Y$ , it should be a conceivable argument that the realistic value of  $\kappa$  cannot be greater or smaller than our present result by a factor of two. That is,  $0.04 \leq \kappa \leq 0.15$  should be a sufficiently generous range of  $\kappa$ . Even if the uncertainty associated with the factorization approximation itself is taken into account, the possibility of  $\kappa \geq 0.2$  would remain extremely small.

A significant deviation of the strong phase difference  $\delta$  from zero, implying significant rescattering effects of  $DK$  states, may reduce the magnitude of  $r$  further. Considering  $DK$  scattering via a  $t$ -channel exchange of Regge trajectories, Deshpande and Dib have made an estimation of the strong phase difference  $(\phi_1 - \phi_0)$  [11]. They obtained  $\tan(\phi_1 - \phi_0) \approx -0.14$ , equivalent to  $\delta \approx 4^\circ$ . This value leads to  $\cos \delta \approx 1$  as an excellent approximation. Nevertheless, one should take such a result more qualitative rather than quantitative, as the method of Regge scattering itself involves large uncertainties. It has been argued that rescattering effects in decays of a  $B$  meson into lighter hadrons might not be as small as commonly imagined [16].

Taking  $\kappa \approx 0.077$ , we obtain the narrow band of  $\gamma$  from Eq. (11):  $87.8^\circ \leq \gamma \leq 92.2^\circ$ . In comparison, analyses of current data show that  $\gamma \sim 65^\circ$  with a generous range  $30^\circ \leq \gamma \leq 150^\circ$  [17]. For illustration we plot  $R_1$  and  $R_2$  as functions of  $\delta$  in Fig. 1, where three different values of  $\gamma$  have been typically taken. One can see that  $R_1 > 1$  and  $R_2 < 1$  for cases (a) and (b); but both  $R_1 > 1$  and  $R_2 > 1$  for case (c), as  $\gamma$  is within the narrow band mentioned above.




Figure 1: Illustrative plot for  $R_{1,2}$ , where  $\kappa = 0.077$  and  $\gamma =$  (a)  $30^\circ$ ; (b)  $60^\circ$ ; and (c)  $90^\circ$ .

It remains unclear that to what extent the constraint  $\sin^2 \gamma \leq R_1 < 1$  or  $\sin^2 \gamma \leq R_2 < 1$  will work.

The idea of extracting the weak phase  $\gamma$  from  $R_{1,2}$  through Eq. (2) or Eq. (12), however, is valid for all allowed values of  $\gamma$ . The feasibility of this method depends on a reliable determination of the coefficient  $\kappa$ . This should be available in the near future, after  $f_D$  is measured from experiments (or calculated from lattice QCD) and the CKM factor  $|V_{ub}/V_{cb}|$  is more accurately extracted from charmless  $B$  decays. At least, our present approach can be complementary to that proposed recently [7] and those suggested previously [2, 3, 5]. It may also confront the nearest data on  $B_u^\pm \rightarrow DK^\pm$  and give a ballpark number to be expected for  $\gamma$ , before the delicate determination of  $\gamma$  becomes available in experiments.

In conclusion, we have shown that it is possible to determine the weak angle  $\gamma$  only from the measurement of the color-allowed  $B_u^\pm \rightarrow DK^\pm$  decay rates. Such measurements are expected to carry out at both  $e^+e^-$   $B$ -meson factories and high-luminosity hadron machines in the coming years.

# References

- [1] CLEO Collaboration, M. Athanas, *et al.*, Report No. hep-ph/9802023.
- [2] M. Gronau and D. Wyler, Phys. Lett. B **265**, 172 (1991).
- [3] M. Gronau and D. London, Phys. Lett. B **253**, 483 (1991); I. Dunietz, Phys. Lett. B **370**, 75 (1991); D. Atwood, G. Eilam, M. Gronau, and A. Soni, Phys. Lett. B **341**, 372 (1995).
- [4] Z.Z. Xing, Nucl. Phys. B (Proc. Suppl.) **50**, 24 (1996); Phys. Lett. B **371**, 310 (1996).
- [5] D. Atwood, I. Dunietz, and A. Soni, Phys. Rev. Lett. **78**, 257 (1997).
- [6] A. Soffer, Report No. CLNS 97/1533 (to be published).
- [7] M. Gronau, Report No. hep-ph/9802315 (to appear in Phys. Rev. Lett.).
- [8] M. Neubert and B. Stech, Report No. hep-ph/9705292.
- [9] M. Bauer, B. Stech, and M. Wirbel, Z. Phys. C **34**, 103 (1987); C **29**, 637 (1985).
- [10] T.E. Browder, K. Honscheid, and D. Pedrini, Ann. Rev. Nucl. Part. Sci. **46**, 395 (1997).
- [11] N.G. Deshpande and C.O. Dib, Phys. Lett. B **319**, 313 (1993).
- [12] Z.Z. Xing, Phys. Rev. D **53**, 2847 (1996).
- [13] B. Blok, M. Gronau, and J.L. Rosner, Phys. Rev. Lett. **78**, 3999 (1997).
- [14] See, e.g., P. Ball, Report No. hep-ph/9802394 (to be published); H.Y. Cheng, C.Y. Cheung, and C.W. Hwang, Phys. Rev. D **55**, 1559 (1997).
- [15] P. Drell, Report No. hep-ex/9711020.
- [16] J.F. Donoghue, E. Golowich, A.A. Petrov, and J.M. Joares, Phys. Rev. Lett. **77**, 2178 (1996).
- [17] F. Parodi, P. Roudeau, and A. Stocchi, Report No. hep-ph/9802289; Y. Nir, Report No. hep-ph/9709301.